

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/370324747>

First records of the blurred lantern shark *Etmopterus bigelowi* from the Cayman Islands, Western Atlantic

Article in *Frontiers in Marine Science* · April 2023

DOI: 10.3389/fmars.2023.1165207

CITATIONS

0

8 authors, including:



Oliver N. Shipley

68 PUBLICATIONS 684 CITATIONS

SEE PROFILE

Some of the authors of this publication are also working on these related projects:



Community dynamics and spatial ecology of elasmobranchs in the New York Bight [View project](#)



Isotopic insights into Bahamian ecosystem dynamics [View project](#)



OPEN ACCESS

EDITED BY
Carlotta Mazzoldi,
University of Padua, Italy

REVIEWED BY
Emilio Sperone,
University of Calabria, Italy
Jorge Paramo,
University of Magdalena, Colombia

*CORRESPONDENCE
Austin J. Gallagher
✉ austin@beneaththewaves.org

SPECIALTY SECTION
This article was submitted to
Discoveries,
a section of the journal
Frontiers in Marine Science

RECEIVED 13 February 2023

ACCEPTED 03 April 2023

PUBLISHED 21 April 2023

CITATION

Gallagher AJ, Shipley ON, De Silva C,
Kohler JK, Fernandes TF, Austin T,
Ormond RF and Gore MA (2023) First
records of the blurred lantern shark
Etmopterus bigelowi from the Cayman
Islands, Western Atlantic.
Front. Mar. Sci. 10:1165207.
doi: 10.3389/fmars.2023.1165207

COPYRIGHT

© 2023 Gallagher, Shipley, De Silva, Kohler,
Fernandes, Austin, Ormond and Gore. This is
an open-access article distributed under the
terms of the [Creative Commons Attribution
License \(CC BY\)](https://creativecommons.org/licenses/by/4.0/). The use, distribution or
reproduction in other forums is permitted,
provided the original author(s) and the
copyright owner(s) are credited and that
the original publication in this journal is
cited, in accordance with accepted
academic practice. No use, distribution or
reproduction is permitted which does not
comply with these terms.

First records of the blurred lantern shark *Etmopterus bigelowi* from the Cayman Islands, Western Atlantic

Austin J. Gallagher^{1*}, Oliver N. Shipley¹, Christine De Silva¹,
Johanna K. Kohler², Teresa F. Fernandes³, Timothy Austin²,
Rupert F. Ormond^{3,4} and Mauvis A. Gore^{3,4}

¹Beneath The Waves, Herndon, VA, United States, ²Cayman Islands Department of Environment, Grand Cayman, Cayman Islands, ³Heriot-Watt University, Edinburgh, United Kingdom, ⁴Marine Conservation International, Edinburgh, United Kingdom

The genus *Etmopterus* is the most speciose group of small bodied deep-sea sharks found throughout the tropical and subtropical Western Atlantic. Despite exhibiting a global distribution at the genus-level, the blurred lantern shark (*Etmopterus bigelowi*) is known only from a few records in the Western and Southern Atlantic Ocean. Through *in-situ* video observations using deep-sea landers, we provide two new locality records of the blurred lantern shark from the deep waters off the Cayman Islands, Caribbean Sea. Three unique individuals were recorded across two separate deployments between 653m – 668m. These observations provide the first records of this species in the Caribbean Sea, adding to the minimal knowledge of the species' distribution throughout the Western Atlantic Ocean.

KEYWORDS

biodiversity, Caribbean, deep-sea, lantern shark, lander, locality, shark

Introduction

Despite their potentially significant contribution to the global biodiversity of fishes, deep-sea sharks remain difficult to observe and study, largely due to the logistical and operational challenges of surveying the deep ocean (Howell et al., 2020). Lantern sharks (Family Etmopteridae) are a small-bodied family of squaliform sharks that are found throughout all major ocean basins and are typically distinguished by light-producing photophores distributed across their body surfaces (Claes and Mallefet, 2011; Claes and Mallefet, 2015). The family includes four extant genera, *Anculeola*, *Centroscyllium*, *Trigonognathus*, and *Etmopterus*, the latter being by far the most speciose, comprising 44 described species (Pollerspöck and Straube, 2022). Species of the genus *Etmopterus* are typically associated with deep-sea environments (*i.e.*, depths exceeding 200m), and as with most species of deep-sea shark, there is an inherent scarcity of basic biological information

pertaining to most aspects of their biology and distribution (e.g., Finucci et al., 2021). This largely reflects the logistical challenges of targeted surveying and the absence of individuals in accessible fish markets, due to their low commercial value (Paramo et al., 2021).

Deep-sea ecosystems of the subtropical and tropical Western Atlantic are believed to support at least eight species of *Etmopterus* (Mejía-Falla et al., 2019). However, most biological information pertaining to these individuals is rudimentary, reflecting morphological and catch data gathered from deep-sea trawl surveys (Springer and Burgess, 1985; Paramo et al., 2012; Paramo et al., 2021). Existing information from waters of the South American Caribbean (Colombia) revealed that catches were largely dominated by a single species, the dwarf lantern shark *E. perryi* (Paramo et al., 2012; Paramo et al., 2021), suggesting comparatively high abundance relative to sympatric species such as *E. robinsoni* and *E. hillianus* (Schofield and Burgess, 1997; Bunkley-Williams and Williams, 2004). In comparison, existing information on *Etmopterus* in the Caribbean is very limited, posing questions on the spatial distribution of species composition throughout the broader regions of the Western Atlantic.

The blurred lantern shark *Etmopterus bigelowi* (Shirai and Tachikawa, 1993) is patchily distributed through insular shelves and slopes of the tropical and subtropical Atlantic, Pacific, and Indian Oceans (Finucci et al., 2020a). Individuals are typically associated with pelagic and benthic deep waters, having been recorded between depths of approximately 100m–1000m (Ebert et al., 2013; Finucci et al., 2020a). Individuals have been found to reach a maximum total length (TL) of at least 73cm; size (TL) at sexual maturity is 40cm – 67cm for males and 50cm – 65cm for females (Ebert et al., 2013). Though populations of *E. bigelowi* are expected to be relatively stable, recent summaries outline a scarcity of information about any aspect of the species' biology (Finucci et al., 2020a). Here, we report the first records of the blurred lantern shark from deep waters adjacent to Grand Cayman, Caribbean Sea, Western Atlantic Ocean, via deployment of deep-sea landers equipped with video cameras. This information provides the first known record of this species in the Caribbean Sea, with previous records only establishing a localized range within South America and the southern United States.

Materials and methods

The Cayman Islands (Grand Cayman, Cayman Brac, and Little Cayman) are situated in the central Caribbean Sea adjacent to the most central portion of Cayman Trench. The region houses a suite of unique oceanographic and geological characteristics due to the proximity of the Oriente transform fault and Cayman Rise, which promote high levels of tectonic activity and hydrothermal vent fields in relatively shallow depths of less than 2,000m (Jones, 1994). Despite the unique oceanographic and marine geological conditions in the region, general trends in biodiversity, especially those related to deep-sea habitats, are poorly described, with the Cayman Islands receiving significantly less attention than similar regions such as the Lucayan Archipelago, Greater Antilles, and Jamaica (Burgess et al., 1994; McLaughlin and Morrissey, 2004; McLaughlin and Morrissey,

2005; Brooks et al., 2015). Further, there have been no systematic survey attempts to quantify patterns of deep-sea biodiversity around the Cayman Islands, despite the potential for housing a suite of unique ecosystems.

As part of a multi-year study to survey the biodiversity of sharks and fishes in the deep-sea territorial waters of the Cayman Islands, we conducted benthic video surveys between March and October 2022 off the shelf in deep waters approximately 3km – 15km offshore spanning the perimeter of Grand Cayman. Surveys were conducted using a custom free-falling, deep baited remote underwater lander system (hereafter dBRUV, Phillips et al., 2019).

The dBRUV system (Figure 1) consisted of a vertically oriented, carbon-fiber frame with pressure-tolerant flotation (G2200, McLane Research Laboratories Inc., East Falmouth, MA) and an acoustic weight-release system (CAT, EdgeTech, Massachusetts, US). This system was equipped with a depth-rated radio transmitter (F1845B, Advanced Telemetry Systems, Minnesota, USA) and an orange identification flag to facilitate recovery at the surface. A single GoPro camera (GoPro Hero 5/6/7), set to record 1080p video at 60 frames per second, was secured within a deep-water housing (GoBenthic, GroupB Incorporated, USA) and attached to the frame ~1.5m above the bottom of the unit. Two LED lights were used to illuminate the seafloor (SiteLite, Juice Robotics, Rhode Island, USA), and the cameras and lights powered with a custom lithium-ion battery pack. Temperature and depth were monitored using a calibrated Starmon TD stand-alone logger (Star Oddi, Iceland) or LAT1400 temperature and depth loggers (LOTEK). Approximately 500g of sardines were attached to a pole-mounted bait cage in front of the camera. The acoustic release system was used to release the drop weight, allowing for the entire system to return to the surface upon command, resulting in deployments of 6–7 hours. All dBRUVs were then located and retrieved at the surface using boat-based GPS unit, YAGI antenna, and VHF radio receiver (R410, Advanced Telemetry Systems, Minnesota, USA). Upon retrieval of the dBRUVs, videos were downloaded from micro-SD cards and reviewed at 0.5x regular speed. All research was conducted under scientific collection permits acquired from the Cayman Islands Department of Environment (DOE).

Results and discussion

On July 26th 2022 a dBRUV was deployed 3km off Southwest Point, Grand Cayman (19.251°N, -81.416°W) at a depth of 653m, which recorded 4.5 hours of video between 10:00 and 16:00. A single male *E. bigelowi* was observed on a single video file and interacted with the bait crate for approximately one minute (Figure 2A). On September 19th, 2022, a dBRUV unit was deployed 15 km south of Breakers Bay, Grand Cayman (19.151°N, -81.191°W) recording 4.5 hours of video at a depth of 668m between 12:00 and 17:00. During the deployment, a single male (Figure 2B) and female (Figure 2C) *E. bigelowi* were observed and interacted with the bait crate for several minutes. Though positive identification of lantern shark species can be inherently challenging (Schofield and Burgess, 1997), the position of the first dorsal fin and snout length distinguishes this




FIGURE 1

Diagram of the deep-sea BRUV (dBRUV) video camera/lander system used in the present study, with its associated components.

species from the sympatric *E. pusillus* (D. Ebert Pers. Comms). These behavioral observations suggest patterns of diel vertical migration in lantern sharks, as seen in other mesopelagic fishes (Abramov and Lipskaya, 1989). As our study suggests, lantern sharks occurring in the Caribbean Sea are likely to be recorded in-between periods of feeding in shallower waters during the night and resting at depth during the day, as documented elsewhere (e.g., Paramo et al., 2021)

These observations support other known localities for *E. bigelowi* from the central Western Atlantic, and represents the

first record of this species from the Caribbean Sea (Table 1). Historical distributions of *E. bigelowi* in Western Atlantic are known from several observations in South America, including Panama, Brazil, Argentina, Suriname, and Uruguay (Table 1), and by a single observation from the USA (Froese and Pauly, 2000). Combined, this suggests that the species displays a moderate latitudinal distribution throughout the tropical and subtropical Western Atlantic, similar to other small and medium-bodied deep-sea sharks such as the Cuban dogfish (*Squalus cubensis*, Cotton et al., 2020) and gulper sharks (*Centrophorus* spp., Finucci




FIGURE 2

In-situ images of *Etmopterus bigelowi* from deep waters of the Cayman Islands collected using deep-sea video landers (Phillips et al., 2019). One male individual was observed on July 26th 2022 (A) followed by observations of a male (B) and female (C) on September 19th 2022. The individuals made several passes across the field of view and were observed interacting with the bait cage.

TABLE 1 Known localities of *Etmopterus bigelowi* throughout Atlantic, Pacific, and Indian Oceans.

Country	Ocean Region	Reference
Liberia	Eastern Atlantic	Froese and Pauly (2023)
South Africa	Eastern Atlantic	Heemstra, 1995
USA	Central Western Atlantic	Froese and Pauly (2023)
Cayman Islands	Central Western Atlantic	This study
Panama	Central Western Atlantic	Froese and Pauly (2023)
Suriname	Central Western Atlantic	Froese and Pauly (2023)
Argentina	Southwestern Atlantic	Menni and Lucifora (2007)
Brazil	Southwestern Atlantic	Soto (2000)
Uruguay	Southwestern Atlantic	Menni and Lucifora (2007)
Hawaii	Central Pacific	Mundy (2005)
Australia	Oceania	Hoese et al. (2006)
Taiwan	Oceania	Froese and Pauly (2022)

et al., 2020b). However, given the very low number of locality records for *E. bigelowi*, it is possible that the species' true latitudinal range remains significantly underestimated.

While the waters of the Cayman Islands are known to support a relatively small, yet diverse population of sharks in its neritic waters (Gore et al., 2020), these new records suggest that its deep waters may have been an overlooked regional hotspot for the biodiversity of deep-sea sharks. As the Cayman Islands have designated 17.8% of their shelf area as Marine Protected Area (MPA) to manage the well-developed and economically important fringing coral reefs, mangrove stands and seagrass beds (Dromard et al., 2011), these MPAs extend only 1-2km offshore. However, the seabed extends to 2000m within Cayman's territorial waters (12 nm), and to over 6000m within Cayman's Exclusive Economic Zone, suggesting that deep-reef habitats may warrant inclusion in future plans for protected area expansion, especially given growing threats of unsustainable, unregulated fishing of deep-water species and of sea-bed mining for metalliferous nodules adjacent to ocean trenches (Irigoin et al., 2014; Churchyard et al., 2016).

These new locality records also highlight a broader need to determine patterns of deep-sea biodiversity throughout the Greater Caribbean (Howell et al., 2021). This is particularly pertinent, given recent calls to scale the size and volume of regional MPAs to facilitate effective conservation of biodiversity (Gallagher et al., 2020). This requires improved efforts to increase region-wide monitoring that will ensure the integration of deep-sea habitats into future MPA expansion (Paramo et al., 2012). To be successful, scaling efforts will require deliberate, international collaboration and improved mechanisms for support and empowerment of many developing countries (Howell et al., 2020; Amon et al., 2022). While providing the critical information required to determine hotspots of deep-sea biodiversity, these efforts are also likely to yield exciting new discoveries across the Greater Caribbean.

Data availability statement

The raw data supporting the conclusions of this article will be made available by the authors, without undue reservation.

Ethics statement

The animal study was reviewed and approved by Heriot-Watt University.

Author contributions

AG, TF, TA and MG led and administered the study. AG, RO, TA and MG conceived the study. All authors performed the field work. OS and JK analyzed the data. OS and AG wrote the first draft of the paper. All authors provided edits on the paper. All authors contributed to the article and approved the submitted version.

Funding

This work was funded through a Darwin Plus Grant (DPLUS140), part of the Darwin Initiative, Overseas Territories Environment and Climate Fund, Department of Environment Food and Rural Affairs of the UK Government, with support from the Department of Environment (DOE) of the Cayman Island Government.

Acknowledgments

We thank staff from the DOE for their assistance with field work and David Ebert for providing key insight on lantern fish biology and positive identification of *E. bigelowi*.

Conflict of interest

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Publisher's note

All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.

References

- Abramov, A., and Lipskaya, H. (1989). On the feeding of *epigonus elegans* in the region of the nazca ridge. *J. Ichthyology* 29, 129–134.
- Amon, D. J., Rotjan, R. D., Kennedy, B. R., Alleng, G., Anta, R., Aram, E., et al. (2022). My deep sea, my backyard: a pilot study to build capacity for global deep-ocean exploration and research. *Philos. Trans. R. Soc. B* 377 (1854), 20210121. doi: 10.1098/rstb.2021.0121
- Brooks, E. J., Brooks, A. M., Williams, S., Jordan, L. K., Abercrombie, D., Chapman, D. D., et al. (2015). First description of deep-water elasmobranch assemblages in the exuma sound, the Bahamas. *Deep Sea Res. Part II: Topical Stud. Oceanography* 115, 81–91.
- Bunkley-Williams, L., and Williams, E. H. (2004). New locality, depth, and size records and species character modifications of some Caribbean deep-reef/shallow slope fishes and a new host and locality record for the chimaera cestodarian. *Caribbean J. Sci.* 40 (1), 88–119.
- Burgess, G. H., Smith, S. H., and Lane, E. D. (1994). “11. fishes of the Cayman islands,” in *The Cayman islands: natural history and biogeography*, (Dordrecht: Kluwer Academic Publishers) vol. 71, 199.
- Churchyard, T., Eaton, M. A., Havery, S., Hall, J., Millett, J., Farr, A., et al. (2016). The biodiversity of the united kingdom’s overseas territories: a stock take of species occurrence and assessment of key knowledge gaps. *Biodiversity Conserv.* 25, 1677–1694. doi: 10.1007/s10531-016-1149-z
- Claes, J. M., and Mallefet, J. (2011). Control of luminescence from lantern shark (*Etmopterus spinax*) photophores. *Communicative Integr. Biol.* 4 (3), 251–253. doi: 10.4161/cib.4.3.14888
- Claes, J. M., and Mallefet, J. (2015). Comparative control of luminescence in sharks: new insights from the slendertail lanternshark (*Etmopterus molleri*). *J. Exp. Mar. Biol. Ecol.* 467, 87–94. doi: 10.1016/j.jembe.2015.03.008
- Cotton, C. F., Derrick, D., Herman, K., Pacoureaux, N., and Dulvy, N. K. (2020). *Squalus cubensis*. The IUCN red list of threatened species 2020. doi: 10.2305/IUCN.UK.2020-3.RLTS.T61416A3104105.en
- Dromard, C. R., Mccoy, C. M., and Turner, J. R. (2011). “Measuring the performance of marine protected areas: the case of little Cayman and Cayman brac, Cayman islands,” in *Proceedings of the 63rd Gulf and Caribbean Fisheries Institute*, San Juan, Puerto Rico. 246–253.
- Ebert, D. A., Fowler, S., and Compagno, L. (2013). *Sharks of the world* (Plymouth: Wild Nature Press).
- Finucci, B., Bineesh, K. K., Cheok, J., Cotton, C. F., Dharmadi, D., Kulka, D. W., et al. (2020b). *Centropristis granulosa*. The IUCN red list of threatened species 2020. doi: 10.2305/IUCN.UK.2020-3.RLTS.T162293947A2897883.en
- Finucci, B., Cheok, J., Cotton, C. F., Kulka, D. W., Neat, F. C., Rigby, C. L., et al. (2020a). *Etmopterus bigelowi*. The IUCN red list of threatened species 2020. doi: 10.2305/IUCN.UK.2020-3.RLTS.T60236A3096618.en
- Finucci, B., Cheok, J., Ebert, D. A., Herman, K., Kyne, P. M., and Dulvy, N. K. (2021). Ghosts of the deep—biodiversity, fisheries, and extinction risk of ghost sharks. *Fish Fisheries* 22 (2), 391–412. doi: 10.1111/faf.12526
- Gallagher, A. J., Amon, D. J., Bervoets, T., Shipley, O. N., Hammerschlag, N., and Sims, D. W. (2020). The Caribbean needs big marine protected areas. *Science* 367 (6479), 749–749.
- Gore, M., Ormond, R., Clarke, C., Kohler, J., Millar, C., and Brooks, E. (2020). Application of photo-identification and lengthened deployment periods to baited remote underwater video stations (BRUVS) abundance estimates of coral reef sharks. *Oceans* 1 (4), 274–299. doi: 10.3390/oceans1040019
- Heemstra, P. C. (1995). “Additions and corrections for the 1995 impression,” in *Revised edition of smiths’ Sea fishes*. Eds. M. M. Smith and P. C. Heemstra (Berlin: Springer-Verlag). p. v-xv.
- Hoesel, D. F., Bray, D. J., Paxton, J. R., and Allen, G. R. (2006). “Fishes,” in *Zoological catalogue of Australia*, vol. Volume 35. Eds. O. L. Beasley and A. Wells (Australia: ABR & CSIRO Publishing), 2178.
- Howell, K. L., Hilário, A., Allcock, A. L., Bailey, D. M., Baker, M., Clark, M. R., et al. (2020). A blueprint for an inclusive, global deep-sea ocean decade field program. *Front. Mar. Sci.* 999. doi: 10.3389/fmars.2020.584861
- Howell, K. L., Hilário, A., Allcock, A. L., Bailey, D., Baker, M., Clark, M. R., et al. (2021). A decade to study deep-sea life. *Nat. Ecol. Evol.* 5 (3), 265–267.
- Irigoin, X., Klevjer, T. A., Røstad, A., Martínez, U., Boyra, G., Acuña, J. L., et al. (2014). Large Mesopelagic fishes biomass and trophic efficiency in the open ocean. *Nat. Commun.* 5 (1), 3271. doi: 10.1038/ncomms4271
- Jones, B. (1994). “Geology of the Cayman islands,” in *The Cayman islands. monographiae biologicae*, vol. vol 71. Eds. M. A. Brunt and J. E. Davies (Dordrecht: Springer). doi: 10.1007/978-94-011-0904-8_2
- McLaughlin, D. M., and Morrissey, J. F. (2004). New records of elasmobranchs from the Cayman trench, Jamaica. *Bull. Mar. Sci.* 75 (3), 481–485.
- McLaughlin, D. M., and Morrissey, J. F. (2005). Reproductive biology of *centrophorus cf. uyato* from the Cayman trench, Jamaica. *J. Mar. Biol. Assoc. United Kingdom* 85 (5), 1185–1192.
- Mejia-Falla, P. A., Navia, A. F., and Salcedo-Reyes, J. C. (2019). Checklist of marine elasmobranchs of Colombia. *Universitas Scientiarum* 24 (1), 164–199.
- Menni, R. C., and Lucifora, L. O. (2007). Condrictios de la Argentina y Uruguay. *ProBiot FcNyM UNLP Serie Técnica-Didáctica La Plata Argent.* 11, 1–15.
- Mundy, B. C. (2005). Checklist of the fishes of the Hawaiian archipelago. *Bishop Mus. Bull. Zool.* 6, 1–704.
- Paramo, J., Wiff, R., and González, R. (2021). A matter of size: the population structure of the smallest known living shark, *Etmopterus perryi* (Springer & burgess 1985), from deep waters off the Colombian Caribbean coast. *J. Fish Biol.* 99 (3), 755–764. doi: 10.1111/jfb.14755
- Paramo, J., Wolff, M., and Saint-Paul, U. (2012). Deep-sea fish assemblages in the Colombian Caribbean Sea. *Fisheries Res.* 125, 87–98. doi: 10.1016/j.fishres.2012.02.011
- Phillips, B. T., Shipley, O. N., Halvorson, J., Sternlicht, J. K., and Gallagher, A. J. (2019). First *in situ* observations of the sharpnose sevengill shark (*Heptranchias perlo*), from the tongue of the ocean, Bahamas. *J. Ocean Sci. Foundation* 32, 17–22.
- Pollerspöck, J., and Straube, N. (2022) *Bibliography database of living/fossil sharks, rays and chimaeras (Chondrichthyes: elasmobranchii, holocephali)*. Available at: <https://shark-references.com/> (Accessed 23 Mar 2023).
- R. Froese and D. Pauly (Eds.) (2000). *FishBase 2000: concepts, design and data sources* (Los Baños, Laguna, Philippines: ICLARM), 344.
- R. Froese and D. Pauly (Eds.) (2022). *FishBase. version (06/2022)*. Available at: <https://www.fishbase.org>.
- R. Froese and D. Pauly (Eds.) (2023). *FishBase. version (06/2023)*. Available at: <https://www.fishbase.org>.
- Schofield, P. J., and Burgess, G. H. (1997). *Etmopterus robinsi* (Elasmobranchii, etmopteridae), a new species of deepwater lantern shark from the Caribbean Sea and western north Atlantic, with a redescription of *Etmopterus hillianus*. *Bull. Mar. Sci.* 60 (3), 1060–1073.
- Shirai, S., and Tachikawa, H. (1993). Taxonomic resolution of the *Etmopterus pusillus* species group (Elasmobranchii, etmopteridae), with description of *E. bigelowi*, n. sp. *Copeia*, 483–495. doi: 10.2307/1447149
- Soto, J. M. R. (2000). Revisão da ocorrência de tubarões etmopteríneos na costa brasileira, com dois novos registros de *Etmopterus bigelowi* shirai & tachikawa 1993 (Chondrichthyes, datatiidae, etmopterinae). *Biotemas* 13 (2), 167–178.
- Springer, S., and Burgess, G. H. (1985). Two new dwarf dogsharks (Etmopterus, squalidae), found off the Caribbean coast of Colombia. *Copeia*, 584–591. doi: 10.2307/1444748